

Review Article

Integrated Use of Soil Ameliorants and Fertilizers to Increase Crop Yield on Acidic Soils of Ethiopian Highlands

Mekonnen Asrat

Department of Plant Sciences, Debre Markos University, Debre Markos, Ethiopia

Email address:

mek.asrat@yahoo.com

To cite this article:Mekonnen Asrat. Integrated Use of Soil Ameliorants and Fertilizers to Increase Crop Yield on Acidic Soils of Ethiopian Highlands. *Advances in Bioscience and Bioengineering*. Vol. 8, No. 2, 2020, pp. 17-23. doi: 10.11648/j.abb.20200802.12**Received:** May 21, 2020; **Accepted:** June 2, 2020; **Published:** July 28, 2020

Abstract: Agriculture has been contributed great share for Ethiopian economy. Ethiopian highlands endowed suitable climatic conditions for crop production, however land degradation in terms of soil acidity and nutrient depletion is becoming the major limiting factor to improve crop yield. Most farmers of Ethiopian highlands are resource poor, small-scale holder, in contrary most crop land characterized by severely eroded and nutrient depleted by interaction effect of environmental conditions and human activities. Around 130 tons/ha soil has been removed from cultivated land, annually besides 40% of the arable land of the country are characterized by acidic soils. To improve crop yield combining use of soil amelioration and fertilization management is critical in such problematic area. Thus, integrated use of 2-3 t/ha lime and NP mineral or organic fertilizers significantly increased crop yield in acidic soils of the country. However, high application rate (11-12 t/ha) of biochar or wood ash on acidic soils has showed equivalent crop yield response to lime rate, hence therefore poor resource farmers can able to ameliorate acidic soils by cheaper and local available materials.

Keywords: Biochare, Fertilizer, Integrated, Lime and Nutrient Depleted

1. Introduction

Ethiopian highlands, areas with altitudes above 1500 masl cover about 500,000 km² it accounts about 45% of the landmass of the country [1]. The area has been contributing 95% of the cultivated land, 90% of the economic activities and home for 88% of the population and 67% livestock of the country [2, 3]. However, the productivity of crop and per capital food production has been declined due to intensive soil fertility depletion in the area [4]. As a result, Ethiopia is frequently affected by food shortages; on average 5 million people require food aid each year. Since 1980s, the country could increase its grain production by 70%, but at the same time total population rose from 40 to 77.1 million and in next 25 years it will further increase by 31 million to 108.7 million [5].

As per [6] report revealed that a series of production failures since the 1970s in Ethiopia have resulted in chronic food insecurity. The causes for food insecurity and poverty may include demographic trends, recurrent drought,

widespread land degradation, shrinking and fragmentation of landholdings, inappropriate policies, poor infrastructure and inefficient agricultural practices [7].

However, soil fertility degradation has been described as the single most important constraint to food security in Sub Saharan Africa (SSA) [8]. Land degradation problem mainly resulting from soil erosion and nutrient depletion, can be singled out as the most important environmental problems creating an unprecedented threat to food security goals of Ethiopia [7]. Most land degradation caused by great pressure from human and livestock populations coupled with other physical political and economical situation of Ethiopian highlands [9]. Most human interference for pursuing economic benefits contributes to rapid and extensive degradation of soils over the past half a century [10]. As per [11] report revealed that 26 percent of the land area in Ethiopia has been degrading over the years 1981-2003, directly affecting the livelihoods of about 29 percent of the population in the country.

Ethiopian agriculture is characterized by smallholder production [12]. One of the major constraints to crop

production faced by smallholder subsistence farmers is inadequate supply of nutrients [13]. Land degradation through soil nutrient depletion in Ethiopia is arising from continuous cropping together with removal of crop residues, low external inputs and absence of adequate soil nutrient saving and recycling technologies [14].

On the other hand, usage of chemical fertilizer in Ethiopia is lower (69.8kg/ha) in terms of application rate per hectare of cultivated land due to various constraints but mainly by economical problem (dominant resource poor farmers) [15]. The use of chemical fertilizers to improve soil fertility of farmland in Ethiopia is low; usually less than 20 kg (N-P) $\text{ha}^{-1} \text{y}^{-1}$ compared to world-wide rate of 97 kg $\text{ha}^{-1} \text{y}^{-1}$, and the fertilizer use is often restricted to irrigated land [16]. As per [17] report indicated that wheat accounted for the largest share (57 kg/ha), and followed by teff (40 k/ha) and maize (29 k/ha), respectively as compared to blanket recommended rate of 200 kg per ha.

Therefore, it is essential to look for agricultural practices that minimize degradation problem and at the same time increase agricultural productivity [7]. Integrated use of soil fertility inputs to alleviating the availability of nutrients problem [18]. On other hand, integrated nutrient management paradigm acknowledges the need for both organic and inorganic material inputs to sustain soil health and crop production due to positive interactions and complements between them [19].

Integrated soil fertility management (ISFM) through provision of optimum fertilizer and soil ameliorants should be coupled with availability of credit facilities contribute to increased access to improved technologies. On the other hand, some recent empirical studies indicated that incorporating use of manure and modest amounts of fertilizers are important to increased and sustained crop yield and becoming more acceptable in smallholder farmers area through considering the economical problem of resource poor farmers [20]. Moreover, the combining use of organic and inorganic nutrient inputs is better option for increasing fertilizer use efficiency and crop yield [21]. The study conducted on *Ferralsols* of Kenya indicated that integrated use of TSP or Minjingu Rock Phosphate (MRP) with


Tithonia significantly increased maize grain yield and alleviate land degradation problem [22].

However, the effectiveness of soil fertility interventions in Ethiopia has historically been constrained by the success in individual programs rather than integrated use of program. Historical evidence in Ethiopia supports the conclusion that the focus on fertilizer to date has been impressive but regrettably has not led to a revolutionary change in yields. Despite a five times increase in fertilizer application, cereal yields have only increased 10 percent since the 1980s [23].

2. Intensity and Valuation of Soil Degradation in Ethiopian Highlands

Soil erosion in agriculture is the most important problem in Ethiopia [24]. Ethiopian highlands soils are severely exposed to water erosion which accounts 130 tons/ha/yr for cropland and 3.4 to 84.5 tons/ha/year soil loosed from all land uses [25] (Figure 1). The problem of accelerating land degradation is especially serious in the intensively cultivated highland parts of the country [26]. Water erosion was the most important process and that in mid 1980's 27 million ha or almost 50% of highland area was significantly eroded, 14 million ha seriously eroded and over 2 million ha beyond reclamation [27].

Soil erosion is estimated to reduce food production by at least 2 percent annually [28]. However, the direct losses of productivity from land degradation are minimally 3 percent of agriculture GDP. With a population growth rate of 2.3 percent this is a critically important [27]. Thus, land degradation has reached a severe stage and has become a major root cause of poverty with significant negative impacts on the country national economy [29]. The direct cost of loss of soil and essential nutrients due to unsustainable land management is estimated to be about three percent of agricultural GDP or \$106 million [27]. As per [30] report estimated that water erosion reduces the potential land for crop production by 10% by 2010 and 30% by 2030 as a result agricultural value per capita drops from \$372 U.S. in 2000 to \$162 in 2030.



Source: Leonard, 2003


Figure 1. The extent and intensity of soil degradation.

The loss of nutrients in Ethiopia is very high more than 60 kg N, P and K ha⁻¹ yr⁻¹ (47 - 60 kg N, -15 kg P₂O₅ and -38 kg K₂O ha⁻¹ [31 32]. On the other hand acid soil infertility is a major limitation to crop production on highly weathered and leached soils of the world. Likewise, about 30% of the highly weathered soils of Ethiopia have been reported to be acidic and it is becoming a serious threat to cereal crops Ethiopian highlands' soils are characterized with low in organic matter content but high in acidity problem due to low land cover [33]. Soil organic matter is derived from decomposed plant and animal residues. Soil acidity is one of the major soil chemical constraints which limit agricultural productivity in the mid and highlands of Ethiopia. Most productive soils have pH in the range of 5.5 to 7.5 while soils with < 5.5 pH categorized in acidic reaction [34]. As per [35] report indicated that 40 percent potential arable land of the country is occupied by acid soils. Out of this, some 15 percent are highly acidic and visible in the west, southern, south-western and northwestern parts of Ethiopia (Figure 2). Soil acidity and its problem are common in all regions where precipitation is high enough to leach appreciable amounts of exchangeable bases from the soil surface. The strongly acid

soils are found in ecology which receive or have historically received high incidence of rainfall and have warm temperatures much of the year [34].

Although acidification is a ongoing natural process in many soil environments, agricultural practices and pollution, mining, and other human activities have accelerated the process [36, 37, 38]. Most Ethiopian highland's soils turned to acidic due to high rainfall and intensive cultivation for long period of time. Highly weathered soils are usually acidic in reaction, low in essential nutrients and organic matter [23, 39]. Moreover, continuous use of acid-forming N fertilizers on highly weathered soils can aggravate acidity problem [40].

Soil acidity produces a number of specific constraints to plant growth. These include elemental toxicities (Al, Mn), reduced availability of nutrients, and reduced rates of certain biological processes and reduced access to soil water. Aluminum toxicity to plants is the main concern farmers with acid soils in Ethiopian highlands [41]. On the other hand, phosphorus ions are adsorbed more strongly on sesquioxide surfaces at lower soil pH due to, in part, the increasing positive charge on these colloids as soil pH decreases [34].



Source: Schleder, 1989

Figure 2. Distribution acidic soils in Ethiopia and Eritrea.

3. Sustainable Soil Fertility Improvement

Ethiopia, an agrarian country whose population and

economy are highly dependent on agriculture, requires to seek ways and means to improve and sustain productivity of its farmland soils through adoption of integrated soil fertility

management practices on a continuous basis with the full participation of farmers [42]. To feed the large and growing population, agricultural production has to be increased by improving the agricultural productivity per land area because most of accessible fertile lands have been cultivated [15]. Ethiopia faces a wider set of soil fertility issues beyond chemical fertilizer use, which has historically been the major focus for extension workers, researchers, policymakers, and donors. If left unchecked, this wider set of issues will limit future output and growth in agriculture across the country; in some areas limit the effectiveness of chemical fertilizer [23].

Integrated plant nutrient management method is need when seeking alternative nutrient management system in

subsistence farming for social and economical acceptability [43]. Fertilization through organic and inorganic sources improves soil fertility of nutrient depleted soils as a result crop production increased. As per [22] report indicated that integrated use of 45 kg/ha TSP and 15 kg/ha Tithonia gave maximum maize grain yield in Kenya. On the other hand, liming is the most effective practice for reducing soil acidity problem [37]. Application of lime provided synergistic effect to TSP and *Tithonia* while slight grain yield increment obtained when TSP combined with lime since P is more limiting than acidity per se while an application of *Tithonia* and lime led increase grain yield by 15 times [22] (it has also shown on Figure 3).




Figure 3. Combine use effect of lime, green manure and P fertilizer on maize grain yield.

Whereas: Treatment 1-control; 2, 4 t/ha lime; 3, Minjingu rock phosphate 60 kg/ha; 4, 60kg/ha TSP; 5, 15 kg/ha Tithonia; 6, 4t/ha lime+ 60 kg/ha rock phosphate; 7, 4t/ha lime+ 60 kg/ha TSP; 8, 4t/ha lime+ 15 kg/ha Tithonia; 9, 45 kg/ha rock phosphate+ 15 kg/ha Tithonia; 10, 45 kg/ha TSP+15 kg/ha Tithonia; 11, 4t/ha lime+ 60 kg/ha rock phosphate+15 kg/ha Tithonia; 12,45 kg/ha TSP+ 15 kg/ha Tithonia+4t/ha lime

Source: Source Tabuet *et al.*, 2007

The integrated use of 32 kg/ha N, 10 kg/ha P and 8 t/ha farmyard produce 3- 3.44 wheat yield (>30% more yield over the control) on medium and low fertile soils of Dila and Dimile [44]. As per [45] study revealed that soil acidity problem can be partly improved by the application of only lime, wheat crop production cannot be improved appreciably without its combined application with NP fertilizers. As in [46] indicated that the combined use of 2 t/ha lime, 40 kg/ha N and 60 kg/ha P₂O₅ fertilizer showed significant effects on teff yield (Table 1).

As per [47] study disclosed that application of wood ash

and manure has significant effect on the properties of acidic soils. In the same study resource soil that received the highest levels of wood ash (11.2 tons CaCO₃ ha⁻¹) showed increment in plant height by 41%, fresh shoot biomass by 100% and dry root biomass by 83%, respectively, compared to the control treatment (Table 2). However, maximum teff grain yield (3.13 t ha⁻¹) was found with combined application of 12 t ha⁻¹ biochar and 100% recommended chemical fertilizer (40/60 N/P₂O₅ kg ha⁻¹) and followed by 2.88 t ha⁻¹ grain yield by using combined 2 t ha⁻¹ lime and 100% recommended chemical fertilizer [46].

Table 1. Combined application of soil ameliorates and chemical fertilizer on tef yield.

Amendments	Fertilizer Rate N/P ₂ O ₅ (kg ha ⁻¹)	Grain yield (t ha ⁻¹)	Dry biomass yield (t ha ⁻¹)
No Amend	0	0.817 ^h	9.22 ^f
	20/30	1.623 ^{ef}	11.54 ^{def}
	40/60	1.870 ^{ef}	13.89 ^{cde}
4 t ha ⁻¹ Biochar	0	0.959 ^h	9.37 ^f
	20/30	1.860 ^f	14.33 ^{cd}
	40/60	2.354 ^{cde}	15.76 ^{bc}
8 t ha ⁻¹ Biochar	0	1.266 ^{gh}	10.40 ^{ef}
	20/30	1.999 ^{def}	13.59 ^{cde}
	40/60	2.676 ^{abc}	17.03 ^{bc}
12 t ha ⁻¹ Biochar	0	2.413 ^{bcd}	16.15 ^{bc}
	20/30	2.462 ^{bcd}	16.14 ^{bc}
	40/60	3.129 ^a	21.04 ^a
2 t ha ⁻¹ Lime	0	2.182 ^{cde}	13.36 ^{cde}
	20/30	2.296 ^{cde}	13.59 ^{cde}
	40/60	2.877 ^{ab}	18.16 ^{ab}
Probability (0.05)		**	**
CV		12.87	12.35

Source: Anteneh et al., 2015

Table 2. Effects of application of lime and wood ash on chemical properties of acidic soils.

Treatment	Rate	pH	Ex. Ac	Ex. Al	CEC	Olsen P	PAIs
			(cmol _c kg ⁻¹)			(mg kg ⁻¹)(%)	
Control	-	4.89 ⁱ	2.22 ^a	1.28 ^a	25.7 ^a	7.0 ^p	5.0 ^a
	3.5	5.15 ^f	0.85 ^f	0.13 ^c	25.8 ^a	7.4 ^{op}	0.5 ^c
Lime (tons CaCO ₃ ha ⁻¹)	7.0	5.43 ^e	0.35 ^h	0.09 ^c	26.1 ^a	7.7 ^{no}	0.3 ^e
	9.2	5.63 ^{bcd}	0.23 ^{ikl}	0.07 ^c	26.4 ^a	8.1 ⁿ	0.3 ^e
	11.2	6.03 ^a	0.14 ^{no}	0.07 ^c	26.4 ^a	8.7 ^m	0.3 ^e
	3.5	5.16 ^{fg}	0.78 ^g	0.12 ^c	25.9 ^a	9.6 ^{kl}	0.5 ^e
Wood ash (tons CaCO ₃ ha ⁻¹)	7.0	5.42 ^c	0.35 ^h	0.10 ^c	26.1 ^a	10.5 ^j	0.4 ^e
	9.2	5.66 ^{bc}	0.19 ^{k-o}	0.08 ^c	26.2 ^a	12.2 ^{gh}	0.3 ^e
	11.2	5.93 ^a	0.16 ^{mno}	0.06 ^c	26.3 ^a	13.8 ^d	0.2 ^e
Mineral P (kg ha ⁻¹) plus lime (7 tons CaCO ₃ ha ⁻¹)	32.5	5.58 ^{cd}	0.27 ^{ij}	0.09 ^c	25.8 ^a	9.0 ^m	0.3 ^e
	65	5.67 ^{bc}	0.23 ^{j-m}	0.08 ^c	25.7 ^a	10.1 ^{ik}	0.3 ^e
	130	5.61 ^{bcd}	0.22 ^{j-m}	0.09 ^c	25.8 ^a	13.1 ^e	0.3 ^e
Manure P (kg ha ⁻¹) plus lime (7 tons CaCO ₃ ha ⁻¹)	32.5	5.57 ^{cd}	0.31 ^{hi}	0.09 ^c	26.4 ^a	8.9 ^m	0.3 ^e
	65	5.64 ^{bcd}	0.20 ^{j-n}	0.10 ^c	26.7 ^a	9.7 ^{kl}	0.4 ^e
	130	5.71 ^b	0.12 ^o	0.09 ^c	26.5 ^a	11.6 ^{hi}	0.3 ^e
Manure P (kg ha ⁻¹) plus wood ash (7 tons CaCO ₃ ha ⁻¹)	32.5	5.62 ^{bcd}	0.32 ^{hi}	0.07 ^c	25.7 ^a	12.5 ^f	0.3 ^e
	65	5.64 ^{bcd}	0.20 ^{ikl}	0.09 ^c	25.8 ^a	14.4 ^c	0.3 ^e
	130	5.62 ^{bcd}	0.16 ^{l-o}	0.07 ^c	26.4 ^a	15.6 ^b	0.3 ^e

Source: Asmare et al., 2015

Moreover, combining use of 5 t/ha manure and 2.2 t/ha lime in northwestern Ethiopian highlands gave maximum bread wheat grain yield (2.7 t/ha) it was 1.98 t/ha more yield over the control treatment [48] (it is located on Table 3).

Table 3. Interaction effect of rate of manure and lime on bread wheat grain yield.

Manure (t ha ⁻¹)	Lime t ha ⁻¹		
	0	2.2	3.3
0	0.898 ^c	1.98 ^b	1.787 ^b
5	1.741 ^b	2.689 ^a	2.169 ^b
10	2.215 ^b	2.736 ^a	2.212 ^{bc}
* Lime and P	1.618 ^c	2.468 ^A	2.056 ^B

Source: Mekonnen et al., 2014

4. Conclusion

Soil nutrient depletion and acidity is the major problem to increase crop production in Ethiopian highlands in area where characterized by subsistence farming system. Integrated nutrient management is the most profitable and achievable method on small-scale holding farming system due to social and economical reasons. Integrated use of lime, chemical and organic fertilizers increased crop yield in Ethiopia highlands, most study findings revealed that combined use of 2-3 t/ha lime, 30-40 kg/ha N and 10-30 kg/ha P chemical fertilizer or 5-8 t/ha manure application improve crop yield in the area. On the other hand, some studies disclosed that lime rate can be substituted by 12 t/ha biochar or 11.2 t/ha wood ash to

ameliorate the problem of acidic soils which is an alternative option for resource poor farmers.

References

- [1] Mohamed-Saleem. Fragile East African highlands: A development vision for smallholder farmers in the Ethiopian highlands. *Outlook on Agriculture*. 1995. 24 (2): 111-116.
- [2] Highlands Reclamation Study. Working paper 17, FAO/Constable, M. Resources of rural development in Ethiopia. Ethiopian Ministry of Agriculture, Addis Ababa. Ethiopia. 1984.
- [3] FAO. Ethiopian Highland Reclamation Study, Ethiopia. Final Report (Volume I and II). Food and Agricultural Organization of the United Nations, Rome, Italy. 1986.
- [4] Amare Hailelassie, Joerg P., Edzo V., Demil Teketay and Jan P. Assessment of soil nutrient depletion and its spatial variability on smallholders' mixed farming systems in Ethiopia using partial versus full nutrient balances. *Elsevier*. www.elsevier.com/locate/agee. *Agriculture, Ecosystems and Environment* 2005, 108, 1–16.
- [5] Haub, C. World population data sheet, Washington, DC; Population Reference Bureau. 2007.
- [6] Kaluski DN, Ophir E, Amede T. Food Security and Nutrition-the Ethiopian Case for Action, *Public Health Nutrition* 2001. 5 (3), 373-381.
- [7] Mengistu Ketema and Siegfried B. Determinants of Manure and Fertilizer Applications in Eastern Highlands of Ethiopia. *Quarterly Journal of International Agriculture*. 2011. 50, No. 3: 237-252.
- [8] World Bank. Country Brief. Development Progress. <http://web.worldbank.org>. 2007.
- [9] Woldeamlak Bewket. Towards integrated watershed management in highland Ethiopia the Chemoga watershed case study. Doctoral Dissertation, Wageningen University, the Netherlands. 2003.
- [10] MEA (Millennium Ecosystem Assessment). Dry land Systems. In *Ecosystem and Well-Being: Current State and Trends*, edited by R. Hassan, R. Schole, and N. Ash, 623–662. Washington, DC, USA. 2005.
- [11] Bai, Z. G., Dent D. L., Olsson L. and Schaepman M. E. Proxy global assessment of land degradation: Review Article. In: *Soil Use and Management*. 2008.24: 223-234.
- [12] AFAP (African Fertilizer and Agribusiness Partnership). Ethiopia Fertilizer Assessment. www.ifdc.org, Alabama, USA. 2012.
- [13] Shapiro, B. I., Sanders, J. H., Fertilizer use in semi-arid West-Africa: profitability and supporting policy. *Agric Syst*. 1998.56: 467-482.
- [14] Sahlemedhin Sertu. Ethiopia: Integrated Soil Management for Sustainable Agriculture and Food Security in Southern and East Africa. Proceedings of the expert consultation, 8-12 December, 1997, Harare, Agri-tex, FAO, 1999. Rome, pp. 197-210.
- [15] Tomoya M. and Takashi Y. The Impacts of Fertilizer Credit on Crop Production and Income in Ethiopia. National Graduate Institute for Policy Studies. 2010.. Tokyo, Japan.
- [16] Pender J, Place F, Ehui S. Strategies for Sustainable Agricultural Development in the Eastern Africa Highlands. EPD Discussion Paper No. 4. IFPRI, Washington, USA. 1999.
- [17] Kefyalew Endale. Fertilizer Consumption and Agricultural Productivity in Ethiopia. Addis Ababa, Ethiopia: EDRI. 2010.
- [18] Palm, C. A., Myers R. J. K. and Nandwa S. M. Combined use of organic and inorganic nutrient sources for soil fertility maintenance and replenishment In: Buresh R. J., Sanchez, P. A., and Calhoun, F. (eds.) *Replenishing Soil Fertility in Africa*. Soil Science Society of America Madison, Wis., 1997. 193-217.
- [19] Sanchez P. A. and Jam B. Soil fertility replenishment takes off in east and southern Africa. In Vanlauwe B. Diels. J Sanginga N. Mercka (eds). *Integrated plant nutrient management in Sub-Saharan Africa. From concept to practice*, CABI, Wallingford UK, 2000. 22-45pp.
- [20] Kimani S. K., Mangale N., Gichuru M., Palm C., Njuho P. and Wamuongo J. Integrated Use and effects of Manure with Modest Application of Inorganic fertilizers on soil Properties and Maize production in the Central Kenya Highlands. Report no. 2 to Rockefellers Foundation, December 1998.
- [21] Mutuo P. K., Mukalama J. P. and Agunda J. On-farm testing of organic and inorganic phosphorous source on maize in Western Kenya. In: *The Biology and Fertility of Tropical Soils: TSBF Report*, 2000. pp 22.
- [22] Tabu I., Bationo A., Obura R. and Masinde J. Effect of Rock Phosphate, Lime and Green Manure on Growth and Yield of Maize in a Non Productive Niche of a Rhodic Ferralsol in Farmer's Fields. In: *Advances in Integrated Soil Fertility Management in sub-Saharan Africa: Challenges and Opportunities*, edited by Andre B., Boaz W. Job K., Joseph K. Springer. 2007.
- [23] IFPRI (International Food Policy Research Institute). Fertilizer and Soil Fertility Potential in Ethiopia, Constraints and Opportunities for Enhancing the System. Sustainable solution for ending hanger and poverty Contributed by Gete Z., Getachew A., Dejene A., Shahid, R. 2010. Working paper, pp 1-66.
- [24] Taffa Tulu. Soil and water conservation, for sustainable agriculture. CTA, Mega publishing. Addis Ababa, Ethiopia. 2002.
- [25] Berry, L. Land degradation in Ethiopia: its extent and impact. Commissioned by the GM with WB support. 2003.
- [26] Woldeamlak Bewket. Soil and Water Conservation Intervention with Conventional Technologies in Northwestern Highlands of Ethiopia: Acceptance and Adoption by Farmers. In: *Land Use Policy*, 2007. 24 (1): 404-416.
- [27] Leonard B., Land Degradation in Ethiopia: its Extent and Impact, Commissioned by the GM with WB support. 2003.
- [28] FAO. 1993. Forest Resource Assessment 1990: Tropical Countries. FAO Forestry Paper Series 112. Rome, Italy.
- [29] Taddese Girma. Land degradation: A challenge to Ethiopia. *Environmental Management*: 2001. 27 (6): 815-824.
- [30] Sonneveld, B. Land Under pressure: The Impact of Water Erosion on Food Production in Ethiopia. Shaker Publishing. Netherlands. 2002.

- [31] Nandwa, S. M. Perspectives on soil fertility in Africa. In: Gichuru, E. K., Ed., *Soil Fertility Management in Africa: A Regional Perspective*, Academy of Science Publishers and Tropical Soil Biology and Fertility Institute of CIAT, Nairobi. 2003.
- [32] Stoorvogel, J. J. and Smaling, E. M. A. 1990. Assessment of soil nutrient depletion in sub-Sahara Africa: 1983-2000. Rep. no. 28. Vol. 1-4. Winand Staring Ctr., Wageningen.
- [33] Tolera Abera., Daba Feyisa and Abebe Zerihun. A Review of Organic and Biological Soil Fertility Management Integrated with NP on Crops Yield and Soil Fertility Improvement in High and Mid Altitude Areas of Western Oromia, Ethiopia. Proceeding on Improved Natural Resource Management Technologies for Food Security, Poverty Reduction and Sustainable Development. Ethiopian Society of Soil Science, March 22-27, EIAR, Addis Ababa, Ethiopia. 2009.
- [34] EIAR (Ethiopian Institute of Agricultural Research). Soil Acidity and Amendments. Miramar, research' in Amharic a monthly newsletter. ISSN 1015-9762, Vol. 9, No. 18. Addis Ababa, Ethiopia. 2009. <http://www.eiar.gov.et>.
- [35] Mesfin Abebe. Nature and Management of Acid Soils in Ethiopia. Addis Ababa, Ethiopia. 2007.
- [36] Curtin D, Syres JK. Lime-induced changes in indices of soil phosphate availability, Soil Science Society of American Journal. 2001. 65, 147-152.
- [37] Fageria, N. K. and Baligar, V. C. Improving nutrient use efficiency of annual crops in Brazilian acid soils for sustainable crop production. Commun. Soil Sci. Plant Anal. 2001. 32: 1303-1319.
- [38] Wang, N. Xu, R. K. and J. Y. Li. Amelioration of an acid Ultisol by agricultural by-products. Land Degradation Development, 2011. 22: 513-518.
- [39] Getachew Agegnehu, Taye Bekele and Agajie Tesfaye. Phosphorus fertilizer and farmyard manure effects on the growth and yield of faba bean and some soil chemical properties in acidic Nitisols of the central highlands of Ethiopia. Ethiopian Journal of Natural Resources, 2005. 7: 23-39.
- [40] Barak, P. Jobe, B. Krueger, A. Peterson, L. and Laird, D. Effects of long-term soil acidification due to nitrogen fertilizer inputs in Wisconsin. Plant and Soil, 197: 61-69 J. Clerk Maxwell, A Treatise on Electricity and Magnetism, 3rd ed., vol. 2. Oxford: Clarendon, 1997, pp. 68-73.
- [41] Fite G, Abdenna D and Wakene N. Utilization of diversity in land use systems: Sustainable and organic approaches to meet human needs. October 2007. Witzhausen, Germany.
- [42] Girmay Gebeyu, Singh B., Mitiku Haila, Borresen T. and Lal R. 2008. Carbon stocks in Ethiopian soils in relation to land use and soil management. Land Degradation & Development 19: 351-367.
- [43] Fageria N. K. and Baligar V. C. 1998. integrated plant nutrient management for sustainable crop production an overview. International journal of tropical agriculture. 15: 1-18.
- [44] Getachew Agegnehu, Ameliorating effects of organic and inorganic fertilizers on crop productivity and soil properties on Reddish-brown soils. In Improved natural resources managements for food security, poverty reduction and sustainable development, Proceeding of the 10th conference of the Ethiopian society of soil science (ESSS). 2009.
- [45] Abreha Kidanemariam. Soil acidity characterization and effects of liming and chemical fertilization on dry matter yield and nutrient uptake of wheat (*Triticum aestivum* L.) on soils of Tsegede District, northern Ethiopia. Doctoral Dissertation, Haramaya University, Haramaya, Ethiopia. 2013.
- [46] Anteneh Abewau, Birru Yitafaru, Yihenew G/Selassie and Tadele Amare. The Role of Biochar on Acid Soil Reclamation and Yield of Teff (*Eragrostis tef* [Zucc] Trotter) in Northwestern Ethiopia. Published by Canadian Center of Science and Education. ISSN 1916-9752 E-ISSN 1916-9760. Journal of Agricultural Science; 2014. Vol. 6, No. 1.
- [47] Asmare M., Markku Y. and Birru Y. Effects of lime, wood Ash, manure and mineral P fertilizer rates on acidity related chemical properties and growth and P uptake of wheat (*Triticum aestivum* L.) on acid soil of Farta district, northwestern highlands of Ethiopia. International Journal of Agriculture and Crop Sciences, SSN 2227-670X. SSN 2227-670X. 2015. Vol 8 (2) 256-269 pp.
- [48] Mekonnen Asrat, Heluf Gebrekidan, Markku, Y. Bobe Bedadi and Wakene Negassa. Effect of integrated use of lime, manure and mineral P fertilizer on bread wheat (*Triticum aestivum* L.) yield, uptake and status of residual soil P on acidic soils of Gozamin District, north-western Ethiopia. Agriculture, Forestry and Fisheries, 2014. 3 (2): 76-85.